

# **Product datasheet for CF502288**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Butyrylcholinesterase (BCHE) Mouse Monoclonal Antibody [Clone ID: OTI4C12]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI4C12

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human BCHE (NP\_000046) produced in HEK293T

cell

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 68.2 kDa

**Gene Name:** butyrylcholinesterase

Database Link: NP 000046

Entrez Gene 590 Human

P06276





### Butyrylcholinesterase (BCHE) Mouse Monoclonal Antibody [Clone ID: OTI4C12] - CF502288

**Background:** Mutant alleles at the BCHE locus are responsible for suxamethonium sensitivity. Homozygous

persons sustain prolonged apnea after administration of the muscle relaxant


suxamethonium in connection with surgical anesthesia. The activity of pseudocholinesterase in the serum is low and its substrate behavior is atypical. In the absence of the relaxant, the

homozygote is at no known disadvantage. [provided by RefSeq]

Synonyms: CHE1; CHE2; E1

**Protein Families:** Druggable Genome, Transmembrane

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BCHE ([RC202198], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BCHE. Positive lysates [LY424950] (100ug) and [LC424950] (20ug) can be purchased separately from OriGene.