

Product datasheet for **CF502650**

DCK Mouse Monoclonal Antibody [Clone ID: OTI16E12]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI16E12
Applications:	FC, WB
Recommended Dilution:	WB 1:2000, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human DCK (NP_000779) produced in HEK293T cell.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	30.3 kDa
Gene Name:	Homo sapiens deoxycytidine kinase (DCK), mRNA.
Database Link:	NP_000779 Entrez Gene 13178 Mouse Entrez Gene 79127 Rat Entrez Gene 1633 Human P27707



[View online »](#)

Background:

Deoxycytidine kinase (DCK) is required for the phosphorylation of several deoxyribonucleosides and their nucleoside analogs. Deficiency of DCK is associated with resistance to antiviral and anticancer chemotherapeutic agents. Conversely, increased deoxycytidine kinase activity is associated with increased activation of these compounds to cytotoxic nucleoside triphosphate derivatives. DCK is clinically important because of its relationship to drug resistance and sensitivity. [provided by RefSeq]

Synonyms:


MGC117410; MGC138632

Protein Families:


Druggable Genome

Protein Pathways:


Purine metabolism, Pyrimidine metabolism

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY DCK ([RC210767], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DCK. Positive lysates [LY400272] (100ug) and [LC400272] (20ug) can be purchased separately from OriGene.



HEK293T cells transfected with either [RC210767] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-DCK antibody ([TA502650]), and then analyzed by flow cytometry.



Flow cytometric Analysis of Jurkat cells, using anti-DCK antibody ([TA502650]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).



Flow cytometric Analysis of HeLa cells, using anti-DCK antibody ([TA502650]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).