

## Product datasheet for **CF502737**

### STAT5A Mouse Monoclonal Antibody [Clone ID: OTI8H1]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI8H1
Applications:	FC, IF, WB
Recommended Dilution:	WB 1:2000, IF 1:100, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human STAT5A (NP_003143) produced in HEK293T cell.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	90.5 kDa
Gene Name:	signal transducer and activator of transcription 5A
Database Link:	<a href="#">NP_003143</a> <a href="#">Entrez Gene 20850 Mouse</a> <a href="#">Entrez Gene 24918 Rat</a> <a href="#">Entrez Gene 6776 Human</a> <a href="#">P42229</a>



[View online »](#)

**Background:**

The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated by, and mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different growth hormones. Activation of this protein in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and has been shown to be essential for the tumorigenesis. The mouse counterpart of this gene is found to induce the expression of BCL2L1/BCL-X(L), which suggests the antiapoptotic function of this gene in cells. [provided by RefSeq]

**Synonyms:**


MGF; STAT5

**Protein Families:**


Druggable Genome, ES Cell Differentiation/IPS, Stem cell relevant signaling - DSL/Notch pathway, Stem cell relevant signaling - JAK/STAT signaling pathway, Transcription Factors

**Protein Pathways:**


Acute myeloid leukemia, Chronic myeloid leukemia, ErbB signaling pathway, Jak-STAT signaling pathway, Pathways in cancer

**Product images:**


HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY STAT5A ([RC205753], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-STAT5A. Positive lysates [LY418866] (100ug) and [LC418866] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (10ug) from 6 different cell lines by using anti-STAT5A monoclonal antibody (1:200).




Anti-STAT5A mouse monoclonal antibody ([TA502737]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY STAT5A ([RC205753]).



HEK293T cells transfected with either [RC205753] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-STAT5A antibody ([TA502737]), and then analyzed by flow cytometry.



Flow cytometric Analysis of HeLa cells, using anti-STAT5A antibody ([TA502737]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-STAT5A antibody ([TA502737]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).