

Product datasheet for CF502849

OriGene Technologies, Inc.9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

PNMT Mouse Monoclonal Antibody [Clone ID: OTI1E8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1E8

Applications: FC, IF, WB

Recommended Dilution: WB 1:2000, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PNMT (NP_002677) produced in HEK293T

cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 30.7 kDa

Gene Name: phenylethanolamine N-methyltransferase

Database Link: NP 002677

Entrez Gene 18948 MouseEntrez Gene 5409 Human

P11086





Background: The product of this gene catalyzes the last step of the catecholamine biosynthesis pathway,

which methylates norepinephrine to form epinephrine (adrenaline). The enzyme also has beta-carboline 2N-methyltransferase activity. This gene is thought to play a key step in


regulating epinephrine production. [provided by RefSeq]

Synonyms: PENT; PNMTase


Protein Families: Druggable Genome

Protein Pathways: Metabolic pathways, Tyrosine metabolism

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PNMT ([RC206586], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PNMT. Positive lysates [LY400946] (100ug) and [LC400946] (20ug) can be purchased separately from OriGene.




Anti-PNMT mouse monoclonal antibody ([TA502849]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PNMT ([RC206586]).





HEK293T cells transfected with either [RC206586] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PNMT antibody ([TA502849]), and then analyzed by flow cytometry.



Flow cytometric Analysis of Jurkat cells, using anti-PNMT antibody ([TA502849]), (Red), compared to a nonspecific negative control antibody, (Blue).