

Product datasheet for CF503332

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NEU2 Mouse Monoclonal Antibody [Clone ID: OTI8C12]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI8C12

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human NEU2(NP_005374) produced in HEK293T

cell

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 42.1 kDa

Gene Name: neuraminidase 2

Database Link: NP 005374

Entrez Gene 4759 Human

Q9Y3R4





NEU2 Mouse Monoclonal Antibody [Clone ID: OTI8C12] - CF503332

Background: This gene belongs to a family of glycohydrolytic enzymes which remove sialic acid residues

from glycoproteins and glycolipids. Expression studies in COS7 cells confirmed that this gene


encodes a functional sialidase. Its cytosolic localization was demonstrated by cell

fractionation experiments. [provided by RefSeq]

Synonyms: SIAL2

Protein Pathways: Other glycan degradation, Sphingolipid metabolism

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NEU2 ([RC219858], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NEU2. Positive lysates [LY401652] (100ug) and [LC401652] (20ug) can be purchased separately from OriGene.