

Product datasheet for CF503341

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

XPF (ERCC4) Mouse Monoclonal Antibody [Clone ID: OTI2C2]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2C2
Applications: FC, IF, WB

Recommended Dilution: WB 1:2000, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ERCC4(NP_005227) produced in HEK293

cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 104.3 kDa

Gene Name: ERCC excision repair 4, endonuclease catalytic subunit

Database Link: NP 005227

Entrez Gene 50505 MouseEntrez Gene 2072 Human

Q92889





Background: The protein encoded by this gene forms a complex with ERCC1 and is involved in the 5'

incision made during nucleotide excision repair. This complex is a structure specific DNA repair endonuclease that interacts with EME1. Defects in this gene are a cause of xeroderma


pigmentosum complementation group F (XP-F), or xeroderma pigmentosum VI (XP6).

Synonyms: ERCC11; FANCQ; RAD1; XFEPS; XPF


Protein Families: Druggable Genome

Protein Pathways: Nucleotide excision repair

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ERCC4 ([RC223300], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ERCC4. Positive lysates [LY401605] (100ug) and [LC401605] (20ug) can be purchased separately from OriGene.



Anti-ERCC4 mouse monoclonal antibody ([TA503341]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ERCC4 ([RC223300]).





HEK293T cells transfected with either [RC223300] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ERCC4 antibody ([TA503341]), and then analyzed by flow cytometry.