

## **Product datasheet for CF503349**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## XPNPEP3 Mouse Monoclonal Antibody [Clone ID: OTI2D5]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2D5

**Applications:** FC, IF, IHC, WB

**Recommended Dilution:** WB 1:500~2000, IHC 1:150, IF 1:50~100, FLOW 1:100

Reactivity: Human, Dog, Rat, Monkey, Mouse

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human XPNPEP3(NP\_071381) produced in HEK293

cell

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 56.9 kDa

**Gene Name:** X-prolyl aminopeptidase 3

Database Link: NP 071381

Entrez Gene 321003 MouseEntrez Gene 685823 RatEntrez Gene 481237 DogEntrez Gene

705778 MonkeyEntrez Gene 63929 Human

Q9NQH7






### Background:

The protein encoded by this gene belongs to the family of X-pro-aminopeptidases that utilize a metal cofactor, and remove the N-terminal amino acid from peptides with a proline residue in the penultimate position. This protein has been shown to localize to the mitochondria of renal cells, and have a role in ciliary function. Mutations in this gene are associated with nephronophthisis-like nephropathy-1. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene, however, expression of some of these isoforms in vivo is not known.


Synonyms: APP3; ICP55; NPHPL1

**Protein Families:** Druggable Genome, Protease

# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY XPNPEP3 (Cat# [RC200888], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-XPNPEP3(Cat# [TA503349]). Positive lysates [LY402903] (100ug) and [LC402903] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-XPNPEP3 monoclonal antibody.






Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-XPNPEP3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA503349])




Anti-XPNPEP3 mouse monoclonal antibody ([TA503349]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY XPNPEP3 ([RC200888]).



Immunofluorescent staining of HeLa cells using anti-XPNPEP3 mouse monoclonal antibody ([TA503349]).





HEK293T cells transfected with either [RC200888] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-XPNPEP3 antibody ([TA503349]), and then analyzed by flow cytometry.