

Product datasheet for CF503486

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MKRN1 Mouse Monoclonal Antibody [Clone ID: OTI3F9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3F9
Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human MKRN1(NP_038474) produced in HEK293T

cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 53.2 kDa

Gene Name: makorin ring finger protein 1

Database Link: NP 038474

Entrez Gene 54484 MouseEntrez Gene 23608 Human

Q9UHC7





MKRN1 Mouse Monoclonal Antibody [Clone ID: OTI3F9] - CF503486

Background: The Makorin ring finger protein-1 gene (MKRN1) is a highly transcribed, intron-containing


source for a family of intronless mammalian genes encoding a novel class of zinc finger proteins. Phylogenetic analyses indicate that the MKRN1 gene is the ancestral founder of this

gene family (Gray et al., 2000 [PubMed 10843807]). [supplied by OMIM]

Synonyms: RNF61

Protein Families: Druggable Genome

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MKRN1 ([RC206848], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MKRN1. Positive lysates [LY415580] (100ug) and [LC415580] (20ug) can be purchased separately from OriGene.