

Product datasheet for CF503677

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GSTO2 Mouse Monoclonal Antibody [Clone ID: OTI4G1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4G1
Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human GSTO2(NP_899062) produced in HEK293T

cell

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 28.1 kDa

Gene Name: glutathione S-transferase omega 2

Database Link: NP 899062

Entrez Gene 309465 RatEntrez Gene 119391 Human

Q9H4Y5

Background: The protein encoded by this gene is an omega class glutathione S-transferase (GST). GSTs are

involved in the metabolism of xenobiotics and carcinogens. Four transcript variants encoding

different isoforms have been found for this gene.






Synonyms: bA127L20.1; GSTO 2-2

Protein Pathways: Drug metabolism - cytochrome P450, Glutathione metabolism, Metabolism of xenobiotics by

cytochrome P450

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY GSTO2 ([RC208961], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GSTO2. Positive lysates [LY405234] (100ug) and [LC405234] (20ug) can be purchased separately from OriGene.