

Product datasheet for CF503975

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PDSS2 Mouse Monoclonal Antibody [Clone ID: OTI1C4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1C4
Applications: IF, WB

Reactivity: WB 1:2000, IF 1:100 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PDSS2(NP_065114) produced in HEK293T

cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 43.9 kDa

Gene Name: decaprenyl diphosphate synthase subunit 2

Database Link: NP 065114

Entrez Gene 71365 MouseEntrez Gene 365592 RatEntrez Gene 57107 Human

Q86YH6






Background:

The protein encoded by this gene is an enzyme that synthesizes the prenyl side-chain of coenzyme Q, or ubiquinone, one of the key elements in the respiratory chain. The gene product catalyzes the formation of all trans-polyprenyl pyrophosphates from isopentyl diphosphate in the assembly of polyisoprenoid side chains, the first step in coenzyme Q biosynthesis. Defects in this gene are a cause of coenzyme Q10 deficiency. COMPLETENESS: complete on the 3' end.


Synonyms: bA59I9.3; C6orf210; COQ10D3; DLP1; hDLP1

Protein Pathways: Terpenoid backbone biosynthesis

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PDSS2 ([RC207892], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDSS2. Positive lysates [LY412512] (100ug) and [LC412512] (20ug) can be purchased separately from OriGene.



Anti-PDSS2 mouse monoclonal antibody ([TA503975]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PDSS2 ([RC207892]).