

Product datasheet for CF504159

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com
CN: techsupport@origene.cn

NLN Mouse Monoclonal Antibody [Clone ID: OTI2F2]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2F2
Applications: FC, WB

Recommended Dilution: WB 1:2000, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human NLN(NP_065777) produced in HEK293T

cell

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 80.5 kDa

Gene Name: neurolysin

Database Link: NP 065777

Entrez Gene 75805 MouseEntrez Gene 117041 RatEntrez Gene 57486 Human

Q9BYT8





Background: This gene encodes a member of the metallopeptidase M3 protein family that cleaves

neurotensin at the Pro10-Tyr11 bond, leading to the formation of neurotensin(1-10) and neurotensin(11-13). The encoded protein is likely involved in the termination of the neurotensinergic signal in the central nervous system and in the gastrointestinal tract.


[provided by RefSeq, Jun

Synonyms: AGTBP; EP24.16; MEP; MOP


Protein Families: Druggable Genome, Protease

Protein Pathways: Renin-angiotensin system

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NLN ([RC212447], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NLN. Positive lysates [LY412353] (100ug) and [LC412353] (20ug) can be purchased separately from OriGene.



Flow cytometric Analysis of Jurkat cells, using anti-NLN antibody ([TA504159]), (Red), compared to a nonspecific negative control antibody, (Blue).