

# **Product datasheet for CF504402**

### **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## **NLN Mouse Monoclonal Antibody [Clone ID: OTI3G8]**

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3G8

Applications: WB

Recommended Dilution: WB 1:500~2000

Reactivity: Human, Rat, Monkey, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human NLN(NP\_065777) produced in HEK293T

cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 80.5 kDa

Gene Name: neurolysin

Database Link: NP 065777

Entrez Gene 75805 MouseEntrez Gene 117041 RatEntrez Gene 696884 MonkeyEntrez Gene

<u>57486 Human</u>

Q9BYT8





**Background:** This gene encodes a member of the metallopeptidase M3 protein family that cleaves


neurotensin at the Pro10-Tyr11 bond, leading to the formation of neurotensin(1-10) and neurotensin(11-13). The encoded protein is likely involved in the termination of the neurotensinergic signal in the central nervous system and in the gastrointestinal tract.

Synonyms: AGTBP; EP24.16; MEP; MOP


**Protein Families:** Druggable Genome, Protease

**Protein Pathways:** Renin-angiotensin system

### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NLN ([RC212447], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NLN. Positive lysates [LY412353] (100ug) and [LC412353] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-NLN monoclonal antibody.