

Product datasheet for CF504472

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CNOT4 Mouse Monoclonal Antibody [Clone ID: OTI3A3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3A3
Applications: WB

"

Recommended Dilution: WB 1:500

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protien fragment corresponding to amino acids 190-455 of human

CNOT4(NP_037448) produced in HEK293T cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 70.2 kDa

Gene Name: CCR4-NOT transcription complex subunit 4

Database Link: NP 037448

Entrez Gene 53621 MouseEntrez Gene 312227 RatEntrez Gene 4850 Human

<u>095628</u>





CNOT4 Mouse Monoclonal Antibody [Clone ID: OTI3A3] - CF504472

Background: The protein encoded by this gene is a subunit of the CCR4-NOT complex, a global

transcriptional regulator. The encoded protein interacts with CNOT1 and has E3 ubiquitin ligase activity. Several transcript variants encoding different isoforms have been found for


this gene.

Synonyms: CLONE243; NOT4; NOT4H

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: RNA degradation

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CNOT4 ([RC217418], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CNOT4. Positive lysates [LY415669] (100ug) and [LC415669] (20ug) can be purchased separately from OriGene.