

Product datasheet for CF504547

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NANP Mouse Monoclonal Antibody [Clone ID: OTI4D5]

Product data:

Isotype:

Product Type: Primary Antibodies

Clone Name: OTI4D5 IHC. WB **Applications:**

Recommended Dilution: WB 1:500, IHC 1:150 Reactivity: Human, Mouse, Rat

Mouse Host: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human NANP(NP_689880) produced in HEK293T

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

> concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Store at -20°C as received. Storage:

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 27.6 kDa

Gene Name: N-acetylneuraminic acid phosphatase

Database Link: NP 689880

Entrez Gene 67311 MouseEntrez Gene 311530 RatEntrez Gene 140838 Human

Q8TBE9


Synonyms: C20orf147; dJ694B14.3; HDHD4

Protein Pathways: Amino sugar and nucleotide sugar metabolism, Metabolic pathways






Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NANP ([RC205503], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NANP. Positive lysates [LY407388] (100ug) and [LC407388] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-NANP mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.