

Product datasheet for CF504982

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

POFUT2 Mouse Monoclonal Antibody [Clone ID: OTI1A6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1A6
Applications: FC, WB

Recommended Dilution: WB 1:1000, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protien of human POFUT2(NP_598368) produced in HEK293T

cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 49.8 kDa

Gene Name: protein O-fucosyltransferase 2

Database Link: NP 598368

Entrez Gene 80294 MouseEntrez Gene 309686 RatEntrez Gene 23275 Human

Q9Y2G5






Background:


Fucose is typically found as a terminal modification of branched chain glycoconjugates, but it also exists in direct O-linkage to serine or threonine residues within cystine knot motifs in epidermal growth factor (EGF; MIM 131530)-like repeats or thrombospondin (THBS; see MIM 188060) type-1 repeats. POFUT2 is an O-fucosyltransferase that use THBS type-1 repeats as substrates (Luo et al., 2006 [PubMed 16464857]). [supplied by OMIM]

Synonyms: C21orf80; FUT13

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY POFUT2 ([RC209505], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-POFUT2. Positive lysates [LY408775] (100ug) and [LC408775] (20ug) can be purchased separately from OriGene.



HEK293T cells transfected with either [RC209505] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-POFUT2 antibody ([TA504982]), and then analyzed by flow cytometry.