

# **Product datasheet for CF505014**

### **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Possible Modern Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MAFB Mouse Monoclonal Antibody [Clone ID: OTI3F6]

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3F6
Applications: FC, WB

Recommended Dilution: WB 1:2000, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 32-320 of human MAFB

(NP 005452) produced in E.coli.

**Formulation:** Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 35.6 kDa

**Gene Name:** MAF bZIP transcription factor B

Database Link: NP 005452

Entrez Gene 16658 MouseEntrez Gene 54264 RatEntrez Gene 9935 Human

Q9Y5Q3





Background: The protein encoded by this gene is a basic leucine zipper (bZIP) transcription factor that


plays an important role in the regulation of lineage-specific hematopoiesis. The encoded nuclear protein represses ETS1-mediated transcription of erythroid-specific genes in myeloid

cells. This gene contains no introns. [provided by RefSeq]


Synonyms: DURS3; KRML; MCTO

**Protein Families:** Druggable Genome, Transcription Factors

## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MAFB ([RC207311], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MAFB. Positive lysates [LY401676] (100ug) and [LC401676] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-MAFB monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human) (1:200).





HEK293T cells transfected with either [RC207311] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-MAFB antibody ([TA505014]), and then analyzed by flow cytometry.