

Product datasheet for CF505015

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

STING (TMEM173) Mouse Monoclonal Antibody [Clone ID: OTI2C9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2C9
Applications: FC, WB

Recommended Dilution: WB 1:200 - 1:1000, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TMEM173(NP_938023) produced in

HEK293T cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 42 kDa

Gene Name: stimulator of interferon response cGAMP interactor 1

Database Link: NP 938023

Entrez Gene 340061 Human

Q86WV6


Synonyms: ERIS; hMITA; hSTING; MITA; MPYS; NET23; SAVI; STING

Protein Pathways: Cytosolic DNA-sensing pathway, RIG-I-like receptor signaling pathway







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TMEM173 ([RC208418], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TMEM173. Positive lysates [LY405006] (100ug) and [LC405006] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 1 cell line by using anti-TMEM173 monoclonal antibody at 1:200.



HEK293T cells transfected with either [RC208418] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-TMEM173 antibody ([TA505015]), and then analyzed by flow cytometry.