

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for CF505205

p15 INK4b (CDKN2B) Mouse Monoclonal Antibody [Clone ID: OTI7C4]

Product data:

Product Type:	Primary Antibodies	
Clone Name:	OTI7C4	
Applications:	WB	
Recommended Dilution:	WB 1:2000	
Reactivity:	Human, Mouse, Rat	
Host:	Mouse	
lsotype:	lgG2b	
Clonality:	Monoclonal	
Immunogen:	Full length human recombinant protein of human CDKN2B(NP_004927) produced in HEK293T cell.	
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)	
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	Unconjugated	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Predicted Protein Size:	14.5 kDa	
Gene Name:	cyclin dependent kinase inhibitor 2B	
Database Link:	<u>NP_004927</u> <u>Entrez Gene 12579 MouseEntrez Gene 25164 RatEntrez Gene 1030 Human</u> <u>P42772</u>	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	p15 INK4b (CDKN2B) Mouse Monoclonal Antibody [Clone ID: OTI7C4] – CF505205
Background:	This gene lies adjacent to the tumor suppressor gene CDKN2A in a region that is frequently mutated and deleted in a wide variety of tumors. This gene encodes a cyclin-dependent kinase inhibitor, which forms a complex with CDK4 or CDK6, and prevents the activation of the CDK kinases, thus the encoded protein functions as a cell growth regulator that controls cell cycle G1 progression. The expression of this gene was found to be dramatically induced by TGF beta, which suggested its role in the TGF beta induced growth inhibition. Two alternatively spliced transcript variants of this gene, which encode distinct proteins, have been reported. [provided by RefSeq]
Synonyms:	CDK4I; INK4B; MTS2; P15; p15INK4b; TP15
Protein Families	S: Druggable Genome
Protein Pathwa	ys: Cell cycle, Pathways in cancer, Small cell lung cancer, TGF-beta signaling pathway

Product images:

~ 火

170	_	
130	_	
100	_	
70	_	
55	_	
40	_	
35	—	
25	—	
15	_	-
10	_	

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CDKN2B ([RC204895], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CDKN2B. Positive lysates [LY417638] (100ug) and [LC417638] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US