

Product datasheet for CF505239

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

UBE2G2 Mouse Monoclonal Antibody [Clone ID: OTI5F1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5F1

Applications: IF, IHC, WB

Recommended Dilution: WB 1:200~2000, IHC 1:150, IF 1:100

Reactivity: Human, Dog, Rat, Monkey, Mouse

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full lenth human recombinant protein of human UBE2G2(NP_003334) produced in HEK293T

cell

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 18.4 kDa

Gene Name: ubiquitin conjugating enzyme E2 G2

Database Link: NP 003334

Entrez Gene 22213 MouseEntrez Gene 294331 RatEntrez Gene 611581 DogEntrez Gene

710862 MonkeyEntrez Gene 7327 Human

P60604





Background:


The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. The encoded protein shares 100% sequence identity with the mouse counterpart. This gene is ubiquitously expressed, with high expression seen in adult muscle. Three alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Jan

Synonyms: UBC7


Protein Families: Druggable Genome

Protein Pathways: Parkinson's disease, Ubiquitin mediated proteolysis

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY UBE2G2 ([RC200407], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-UBE2G2. Positive lysates [LY418761] (100ug) and [LC418761] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by usin g anti-UBE2G2 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-UBE2G2 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-UBE2G2 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-UBE2G2 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-UBE2G2 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Anti-UBE2G2 mouse monoclonal antibody ([TA505239]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY UBE2G2 ([RC200407]).