

# **Product datasheet for CF505416**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### HTF9C (TRMT2A) Mouse Monoclonal Antibody [Clone ID: OTI2A1]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2A1

Applications: IHC, WB

**Reactivity:** WB 1:2000, IHC 1:150 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TRMT2A(NP\_073564) produced in HEK293T

cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 68.5 kDa

**Gene Name:** tRNA methyltransferase 2 homolog A

Database Link: NP 073564

Entrez Gene 27037 Human

Q8IZ69






Background:


The protein encoded by this gene is of unknown function. However, it is orthologous to the mouse Trmt2a gene and contains an RNA methyltransferase domain. Expression of this gene varies during the cell cycle, with aberrant expression being a possible biomarker in certain breast cancers. Several transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Apr 2012]


Synonyms: HTF9C

**Protein Families:** Druggable Genome

## **Product images:**









Equivalent amounts of cell lysates (10 ug per lane) of wild-type HeLa cells (WT, Cat# LC810HELA) and TRMT2A-Knockout HeLa cells (KO, Cat# [LC833845]) were separated by SDS-PAGE and immunoblotted with anti-TRMT2A monoclonal antibody [TA505416] (1:500). Then the blotted membrane was stripped and reprobed with anti-PCNA antibody as a loading control.





Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-TRMT2A mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-TRMT2A mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.