

Product datasheet for CF505491

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GADD45G Mouse Monoclonal Antibody [Clone ID: OTI1G2]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1G2
Applications: FC, IF, WB

Recommended Dilution: WB 1:4000, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human GADD45G(NP_006696) produced in

HEK293T cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 16.9 kDa

Gene Name: growth arrest and DNA damage inducible gamma

Database Link: NP 006696

Entrez Gene 23882 MouseEntrez Gene 291005 RatEntrez Gene 10912 Human

O95257



GADD45G Mouse Monoclonal Antibody [Clone ID: OTI1G2] - CF505491

Background: This gene is a member of a group of genes whose transcript levels are increased following


stressful growth arrest conditions and treatment with DNA-damaging agents. The protein encoded by this gene responds to environmental stresses by mediating activation of the p38/JNK pathway via MTK1/MEKK4 kinase. The GADD45G is highly expressed in placenta.

[provided by RefSeq]


Synonyms: CR6; DDIT2; GADD45gamma; GRP17

Protein Pathways: Cell cycle, MAPK signaling pathway, p53 signaling pathway

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY GADD45G ([RC201364], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GADD45G. Positive lysates [LY416476] (100ug) and [LC416476] (20ug) can be purchased separately from OriGene.



Anti-GADD45G mouse monoclonal antibody ([TA505491]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY GADD45G ([RC201364]).





HEK293T cells transfected with either [RC201364] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-GADD45G antibody ([TA505491]), and then analyzed by flow cytometry.