

Product datasheet for CF505561

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TRIM44 Mouse Monoclonal Antibody [Clone ID: OTI1E10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1E10

Applications: IF, IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150, IF 1:100

Reactivity: Human, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TRIM44(NP_060053) produced in HEK293T

cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 38.3 kDa

Gene Name: tripartite motif containing 44

Database Link: NP 060053

Entrez Gene 54765 Human

Q96DX7

Background: This gene encodes a member of the tripartite motif (TRIM) family. The TRIM motif includes

three zinc-binding domains, namely a RING, a B-box type 1 and a B-box type 2, and a coiled-

coil region. [provided by RefSeq]







Synonyms:

DIPB; HSA249128; MC7


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY TRIM44 (Cat# [RC201074], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TRIM44 (Cat# [TA505561])(1:500)




Western blot analysis of extracts (35ug) from 6 cell line lysates by using anti-TRIM44 monoclonal antibody (1:500).



Immunohistochemical staining of paraffinembedded Human breast tissue within the normal limits using anti-TRIM44 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-TRIM44 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-TRIM44 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.







Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-TRIM44 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-TRIM44 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-TRIM44 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-TRIM44 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-TRIM44 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-TRIM44 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Anti-TRIM44 mouse monoclonal antibody ([TA505561]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY TRIM44 ([RC201074]).