

### **Product datasheet for CF505609**

# OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## **POGK Mouse Monoclonal Antibody [Clone ID: OTI2G8]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2G8
Applications: IHC, WB

Recommended Dilution: WB 1:500, IHC 1:150, IF 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human POGK(NP\_060012) produced in HEK293T

cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 69.3 kDa

**Gene Name:** pogo transposable element derived with KRAB domain

Database Link: NP 060012

Entrez Gene 57645 Human

Q9P215






Background:


The exact function of the protein encoded by this gene is not known. However, this gene product contains a KRAB domain (which is involved in protein-protein interactions) at the N-terminus, and a transposase domain at the C-terminus, suggesting that it may belong to the family of DNA-mediated transposons in human. [provided by RefSeq, Jul 2008]

Synonyms: BASS2; KRBOX2; LST003 **Protein Families:** Transcription Factors

#### **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY POGK ([RC222119], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-POGK. Positive lysates [LY413730] (100ug) and [LC413730] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human breast tissue within the normal limits using anti-POGK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505609])






Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-POGK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505609])




Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-POGK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505609])




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-POGK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505609])






Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-POGK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505609])




Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-POGK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505609])



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-POGK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505609])





Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-POGK mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA505609])