

Product datasheet for CF506345

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD64 (FCGR1A) Mouse Monoclonal Antibody [Clone ID: OTI3H9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3H9
Applications: FC, IF, WB

Recommended Dilution: WB 1:1000, IF 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human FCGR1A(NP_000557) produced in HEK293T

cell

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 42.5 kDa

Gene Name: Fc fragment of lgG receptor la

Database Link: NP 000557

Entrez Gene 2209 Human

P12314

Background: This gene encodes a protein that plays an important role in the immune response. This

protein is a high-affinity Fc-gamma receptor. The gene is one of three related gene family

members located on chromosome 1. [provided by RefSeq, Jul 2008]






Synonyms: CD64; CD64A; FCRI; IGFR1

Protein Families: Transmembrane


Protein Pathways: Fc gamma R-mediated phagocytosis, Hematopoietic cell lineage, Systemic lupus

erythematosus


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY FCGR1A ([RC207487], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FCGR1A. Positive lysates [LY400193] (100ug) and [LC400193] (20ug) can be purchased separately from OriGene.




Immunofluorescent staining of HeLa cells using anti-FCGR1A mouse monoclonal antibody ([TA506345]).



Anti-FCGR1A mouse monoclonal antibody ([TA506345]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY FCGR1A ([RC207487]).





Flow cytometric analysis of living 293T cells transfected with FCGR1A overexpression plasmid ([RC207487]), Red)/empty vector ([PS100001], Blue) using anti-FCGR1A antibody ([TA506345]). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).