

Product datasheet for CF506883

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Methylmalonyl Coenzyme A mutase (MUT) Mouse Monoclonal Antibody [Clone ID: OTI2A8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2A8
Applications: IF, WB

Recommended Dilution: WB 1:1000 -1:4000, IF 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human MUT(NP_000246) produced in HEK293T

cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 79.3 kDa

Gene Name: methylmalonyl-CoA mutase

Database Link: NP 000246

Entrez Gene 4594 Human

P22033





Methylmalonyl Coenzyme A mutase (MUT) Mouse Monoclonal Antibody [Clone ID: OTI2A8] – CF506883

Background: This gene encodes the mitochondrial enzyme methylmalonyl Coenzyme A mutase. In

humans, the product of this gene is a vitamin B12-dependent enzyme which catalyzes the isomerization of methylmalonyl-CoA to succinyl-CoA, while in other species this enzyme may have different functions. Mutations in this gene may lead to various types of methylmalonic


aciduria. [provided by RefSeq, Jul 2008]

Synonyms: MCM


Protein Families: Druggable Genome

Protein Pathways: Metabolic pathways, Propanoate metabolism, Valine, leucine and isoleucine degradation

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MUT ([RC201972], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MUT. Positive lysates [LY400099] (100ug) and [LC400099] (20ug) can be purchased separately from OriGene.



Immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY MUT ([RC201972]) using Anti-MUT mouse monoclonal antibody ([TA506883]) at 1:100 dilution.