

# **Product datasheet for CF800425**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **ZSCAN4 Mouse Monoclonal Antibody [Clone ID: OTI1E7]**

Human

#### **Product data:**

Reactivity:

**Product Type:** Primary Antibodies

Clone Name: OTI1E7
Applications: WB

Applications.

**Recommended Dilution:** WB: 1:2000

Host: Mouse

Isotype: lgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 164-433 of human

ZSCAN4 (NP 689890) produced in E.coli.

**Formulation:** Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 48.8 kDa

**Gene Name:** zinc finger and SCAN domain containing 4

Database Link: NP 689890

Entrez Gene 201516 Human

Q8NAM6





## ZSCAN4 Mouse Monoclonal Antibody [Clone ID: OTI1E7] - CF800425

**Background:** The ZSCAN4 gene encodes a protein involved in telomere maintenance and with a key role in


the critical feature of mouse embryonic stem (ES) cells, namely, defying cellular senescence and maintaining normal karyotype for many cell divisions in culture (Zalzman et al., 2010

[PubMed 20336070]). [supplied by OMIM]

Synonyms: ZNF494

**Protein Families:** Transcription Factors

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ZSCAN4 ([RC211003], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ZSCAN4. Positive lysates [LY407355] (100ug) and [LC407355] (20ug) can be purchased separately from OriGene.