

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for CF800609

HBS1L Mouse Monoclonal Antibody [Clone ID: OTI1D5]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1D5
Applications:	WB
Recommended Dilution:	WB 1:2000
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 1-260 of human HBS1L (NP_006611) produced in E.coli.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	75.3 kDa
Gene Name:	HBS1 like translational GTPase
Database Link:	<u>NP_006611</u> <u>Entrez Gene 56422 MouseEntrez Gene 293408 RatEntrez Gene 10767 Human</u> <u>Q9Y450</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE HBS1L Mouse Monoclonal Antibody [Clone ID: OTI1D5] – CF800609

Background:This gene encodes a member of the GTP-binding elongation factor family. It is expressed in
multiple tissues with the highest expression in heart and skeletal muscle. The intergenic
region of this gene and the MYB gene has been identified to be a quantitative trait locus
(QTL) controlling fetal hemoglobin level, and this region influnces erythrocyte, platelet, and
monocyte counts as well as erythrocyte volume and hemoglobin content. DNA
polymorphisms at this region associate with fetal hemoglobin levels and pain crises in sickle
cell disease. A single nucleotide polymorphism in exon 1 of this gene is significantly
associated with severity in beta-thalassemia/Hemoglobin E. Multiple alternatively spliced
transcript variants encoding different protein isoforms have been found for this gene.
[provided by RefSeq]

Synonyms:

EF-1a; eRF3c; ERFS; HBS1; HSPC276

Product images:


158-106-

79-

48-

35-

23-



HepG2 HeLa SVT2 A549 COS7 Jurkat MDCK PC12 MCF7

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HBS1L ([RC208125], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HBS1L. Positive lysates [LY416509] (100ug) and [LC416509] (20ug) can be purchased separately from OriGene.

Western blot analysis of extracts (35ug) from 9 different cell lines by usin g anti-HBS1L monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

