

Product datasheet for **CF801631**

Dicer (DICER1) Mouse Monoclonal Antibody [Clone ID: OT1E11]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OT1E11
Applications:	IHC
Recommended Dilution:	IHC 1:150
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 1666-1922 of human DICER1 (NP_085124) produced in E.coli.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	218.5 kDa
Gene Name:	dicer 1, ribonuclease III
Database Link:	NP_085124 Entrez Gene 192119 Mouse Entrez Gene 299284 Rat Entrez Gene 23405 Human Q9UPY3




[View online »](#)

Background: This gene encodes a protein possessing an RNA helicase motif containing a DEXH box in its amino terminus and an RNA motif in the carboxy terminus. The encoded protein functions as a ribonuclease and is required by the RNA interference and small temporal RNA (stRNA) pathways to produce the active small RNA component that represses gene expression. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep


Synonyms: DCR1; Dicer; Dicer1e; HERNA; K12H4.8-LIKE; MNG1; RMSE2

Protein Families: Druggable Genome

Product images:



Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-DICER1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA801631])



Immunohistochemical staining of paraffin-embedded Human lymph node tissue within the normal limits using anti-DICER1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA801631])