

## **Product datasheet for CF801840**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## HMG2L1 (HMGXB4) Mouse Monoclonal Antibody [Clone ID: OTI1H9]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1H9

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 1-348 of human

HMGXB4 (NP\_005478) produced in E.coli.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 53 kDa

Gene Name: HMG-box containing 4

Database Link: NP 005478

Entrez Gene 10042 Human

Q9UGU5

Background: High mobility group (HMG) proteins are nonhistone chromosomal proteins. See HMG2 (MIM

163906) for additional information on HMG proteins. [supplied by OMIM, Nov 2010]






### HMG2L1 (HMGXB4) Mouse Monoclonal Antibody [Clone ID: OTI1H9] - CF801840

Synonyms: high-mobility group protein 2-like 1; HMG2L1; HMGBCG; HMG box domain containing 4;

OTTHUMP00000028778; THC211630

**Protein Families:** Transcription Factors

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HMGXB4 ([RC211137], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HMGXB4. Positive lysates [LY417272] (100ug) and [LC417272] (20ug) can be purchased separately from OriGene.