

Product datasheet for CF802412

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

EU: info-de@origene.com CN: techsupport@origene.cn

CD5 Mouse Monoclonal Antibody [Clone ID: OTI3E5]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3E5
Applications: FC, WB
Recommended Dilution: WB 1:500
Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CD5 (NP_055022) produced in HEK293T

cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 54.4 kDa

Gene Name: CD5 molecule

Database Link: NP 055022

Entrez Gene 921 Human

P06127

Synonyms: LEU1; T1


Protein Families: Druggable Genome, Transmembrane






Protein Pathways: Hematopoietic cell lineage

Product images:





HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CD5 ([RC206494], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD5. (1:5. Positive lysates [LY415443] (100ug) and [LC415443] (20ug) can be purchased separately from OriGene.




Living HEK293T cells transfected with either [RC206494] overexpress plasmid (Red), compared to an IgG isotype control, (Green) or empty vector control plasmid (Blue) were immunostained by anti-CD5 antibody ([TA802412]), and then analyzed by flow cytometry (1:100).









Flow cytometric analysis of living CCRF-CEM cells, using anti-CD5 antibody ([TA802412]), (red), compared to an IgG isotype control, (green), and negative control (PBS), (blue) (1:100).