

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for CF804016

CD63 Mouse Monoclonal Antibody [Clone ID: OTI3G6]

Product data:

Product Type:	Primary Antibodies	
Clone Name:	OTI3G6	
Applications:	FC	
Recommended Dilution:	FLOW 1:50	
Reactivity:	Human	
Host:	Mouse	
lsotype:	lgG1	
Clonality:	Monoclonal	
Immunogen:	Full length human recombinant protein of human CD63 (NP_001771) produced in HEK293T cell.	
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)	
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	Unconjugated	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Predicted Protein Size:	25.5 kDa	
Gene Name:	CD63 molecule	
Database Link:	<u>NP_001771</u> <u>Entrez Gene 967 Human</u> <u>P08962</u>	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


CD63 Mouse Monoclonal Antibody [Clone ID: OTI3G6] – CF804016

characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. The encoded protein is a cell surface glycoprotein that is known to complex with integrins. It may function as a blood platelet activation marker. Deficiency of this protein is associated with Hermansky-Pudlak syndrome. Also this gene has been associated with tumor progression. Alternative splicing results in multiple transcript variants encoding different protein isoforms. [provided by RefSeq, Apr 2012]	Background:	transduction events that play a role in the regulation of cell development, activation, growth and motility. The encoded protein is a cell surface glycoprotein that is known to complex wit integrins. It may function as a blood platelet activation marker. Deficiency of this protein is associated with Hermansky-Pudlak syndrome. Also this gene has been associated with tumo progression. Alternative splicing results in multiple transcript variants encoding different
---	-------------	---

Synonyms:	LAMP-3; ME491; MLA1; OMA81H; TSPAN30
Protein Families:	Druggable Genome, Transmembrane

Protein Pathways: Lysosome

Product images:



Flow cytometric Analysis of living A549 cells, using anti-CD63 antibody ([TA804016]), (Red), compared to a nonspecific negative control antibody, (Blue).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US