

### **Product datasheet for CF805450**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## hSET1 (SETD1A) Mouse Monoclonal Antibody [Clone ID: OTI1D3]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1D3
Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 1-320 of human

SETD1A(NP\_055527) produced in E.coli.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 185.9 kDa

**Gene Name:** SET domain containing 1A, histone lysine methyltransferase

Database Link: NP 055527

Entrez Gene 233904 MouseEntrez Gene 9739 Human

015047





### hSET1 (SETD1A) Mouse Monoclonal Antibody [Clone ID: OTI1D3] - CF805450

**Background:** SET1A is a component of a histone methyltransferase (HMT) complex that produces mono-,

di-, and trimethylated histone H3 at Lys4. The complex is the analog of the S. cerevisiae Set1/COMPASS complex (Lee and Skalnik, 2005 [PubMed 16253997]). Also see SET1B (MIM


611055). [supplied by OMIM, Mar 2008]

Synonyms: KMT2F; Set1; Set1A

Protein Families: Druggable Genome

Protein Pathways: Lysine degradation

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SETD1A ([RC214996], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SETD1A. Positive lysates [LY415047] (100ug) and [LC415047] (20ug) can be purchased separately from OriGene.