

## Product datasheet for **CF806459**

### Retinoic Acid Receptor alpha (RARA) Mouse Monoclonal Antibody [Clone ID: OTI7B8]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI7B8
Applications:	WB
Recommended Dilution:	WB 1:2000
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 52-299 of human RARA(NP_000955) produced in E.coli.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	50.6 kDa
Gene Name:	retinoic acid receptor alpha
Database Link:	<a href="#">NP_000955</a> <a href="#">Entrez Gene 19401 Mouse</a> <a href="#">Entrez Gene 24705 Rat</a> <a href="#">Entrez Gene 5914 Human</a> <a href="#">P10276</a>



[View online »](#)

**Background:**

This gene represents a nuclear retinoic acid receptor. The encoded protein, retinoic acid receptor alpha, regulates transcription in a ligand-dependent manner. This gene has been implicated in regulation of development, differentiation, apoptosis, granulopoiesis, and transcription of clock genes. Translocations between this locus and several other loci have been associated with acute promyelocytic leukemia. Alternatively spliced transcript variants have been found for this locus. [provided by RefSeq, Sep 2010]

**Synonyms:**


NR1B1; RAR

**Protein Families:**

Druggable Genome, Nuclear Hormone Receptor, Transcription Factors

**Protein Pathways:**

Acute myeloid leukemia, Pathways in cancer

**Product images:**

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY RARA ([RC200430], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RARA. Positive lysates [LY400350] (100ug) and [LC400350] (20ug) can be purchased separately from OriGene.