

### **Product datasheet for CF806584**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## HNF 4 alpha (HNF4A) Mouse Monoclonal Antibody [Clone ID: OTI5A9]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI5A9

**Applications:** WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 1-315 of human

HNF4A(NP\_849180) produced in E.coli.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** Homo sapiens hepatocyte nuclear factor 4 alpha (HNF4A), transcript variant 1, mRNA.

Database Link: NP 849180

Entrez Gene 15378 MouseEntrez Gene 25735 RatEntrez Gene 3172 Human

P41235

Synonyms: FRTS4; HNF4; HNF4a7; HNF4a8; HNF4a9; HNF4alpha; MODY; MODY1; NR2A1; NR2A21; TCF;

TCF-14; TCF14

**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Nuclear Hormone Receptor, Transcription

**Factors** 






**Protein Pathways:** 

Maturity onset diabetes of the young

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HNF4A ([RC214914], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HNF4A.