

## **Product datasheet for CF806894**

#### **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

# FLT4 Mouse Monoclonal Antibody [Clone ID: OTI1G7]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1G7
Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 1063-1363 of human

FLT4(NP 891555) produced in E.coli.

**Formulation:** Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 150.2 kDa

**Gene Name:** fms related receptor tyrosine kinase 4

Database Link: NP 891555

Entrez Gene 14257 MouseEntrez Gene 114110 RatEntrez Gene 2324 Human

<u>P35916</u>





### FLT4 Mouse Monoclonal Antibody [Clone ID: OTI1G7] - CF806894

**Background:** This gene encodes a tyrosine kinase receptor for vascular endothelial growth factors C and D.


The protein is thought to be involved in lymphangiogenesis and maintenance of the lymphatic endothelium. Mutations in this gene cause hereditary lymphedema type IA.

[provided by RefSeq, Jul 2008]

Synonyms: FLT-4; FLT41; LMPH1A; PCL; VEGFR-3; VEGFR3

Protein Families: Druggable Genome, Protein Kinase, Transmembrane
Protein Pathways: Cytokine-cytokine receptor interaction, Focal adhesion

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY FLT4 ([RC214285], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FLT4. Positive lysates [LY405313] (100ug) and [LC405313] (20ug) can be purchased separately from OriGene.