

Product datasheet for CF807197

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD45 (PTPRC) Mouse Monoclonal Antibody [Clone ID: UCH-L1]

Product data:

Product Type: Primary Antibodies

Clone Name: UCH-L1
Applications: FC, IHC
Recommended Dilution: IHC 1:150

Reactivity:HumanHost:MouseIsotype:IgG2a

Clonality: Monoclonal

Immunogen: IL-2-dependent T-cell line (CA1).

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: protein tyrosine phosphatase receptor type C

Database Link: NP 563578

Entrez Gene 5788 Human

P08575





Background:


The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitosis, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported. [provided by RefSeq, Jun 2012]

Synonyms: B220; CD45; CD45R; GP180; L-CA; LCA; LY5; T200

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Phosphatase, Transmembrane


Protein Pathways: Cell adhesion molecules (CAMs), Fc gamma R-mediated phagocytosis, Primary immunodeficiency, T cell receptor signaling pathway

Product images:




Immunohistochemical staining of paraffinembedded Human liver tissue using anti-CD45RO mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.






Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-CD45RO mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-CD45RO mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-CD45RO mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.







Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-CD45RO mouse monoclonal antibody. Heatinduced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Human tonsil using anti-CD45RO mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Flow cytometric Analysis of living RBC-lysed human peripheral blood cells, using anti-CD45RO antibody ([TA807197]), (Red), compared to a nonspecific negative control antibody ([TA180143]), (green) and PBS, (blue) (1:100).



Flow cytometric Analysis of living Jurkat cells, using anti-CD45RO antibody ([TA807197]), (Red), compared to a nonspecific negative control antibody ([TA180143]), (green), and PBS, (blue) (1:100).