

## Product datasheet for CF807259

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **HIPK1 Mouse Monoclonal Antibody [Clone ID: OTI4B7]**

WB 1:2000

#### **Product data:**

**Product Type: Primary Antibodies** 

Clone Name: OTI4B7

**Applications:** 

Recommended Dilution:

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 439-697 of human

HIPK1(NP 852003) produced in E.coli.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

> concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Store at -20°C as received. Storage:

Stability: Stable for 12 months from date of receipt.

**Predicted Protein Size:** 87.1 kDa

Gene Name: homeodomain interacting protein kinase 1

Database Link: NP 852003

Entrez Gene 15257 MouseEntrez Gene 365895 RatEntrez Gene 204851 Human

Q86Z02






**Background:** The protein encoded by this gene belongs to the Ser/Thr family of protein kinases and HIPK

subfamily. It phosphorylates homeodomain transcription factors and may also function as a co-repressor for homeodomain transcription factors. Alternative splicing results in four transcript variants encoding four distinct isoforms. [provided by RefSeq, Jul 2008]

Synonyms: Myak; Nbak2

**Protein Families:** Druggable Genome, Protein Kinase, Transcription Factors

### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HIPK1 ([RC206176], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HIPK1 (1:2000). Positive lysates [LY405750] (100ug) and [LC405750] (20ug) can be purchased separately from OriGene.