

## **Product datasheet for CF807822**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Troponin I fast skeletal muscle (TNNI2) Mouse Monoclonal Antibody [Clone ID: OTI1H1]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1H1
Applications: WB

rippineacions.

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human TNNI2 (NP\_003273) produced in E.coli.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 21.2 kDa

Gene Name: troponin I2, fast skeletal type

Database Link: NP 003273

Entrez Gene 21953 MouseEntrez Gene 29389 RatEntrez Gene 7136 Human

P48788



# Troponin I fast skeletal muscle (TNNI2) Mouse Monoclonal Antibody [Clone ID: OTI1H1] – CF807822


Background:

This gene encodes a fast-twitch skeletal muscle protein, a member of the troponin I gene family, and a component of the troponin complex including troponin T, troponin C and troponin I subunits. The troponin complex, along with tropomyosin, is responsible for the calcium-dependent regulation of striated muscle contraction. Mouse studies show that this component is also present in vascular smooth muscle and may play a role in regulation of smooth muscle function. In addition to muscle tissues, this protein is found in corneal epithelium, cartilage where it is an inhibitor of angiogenesis to inhibit tumor growth and metastasis, and mammary gland where it functions as a co-activator of estrogen receptor-related receptor alpha. This protein also suppresses tumor growth in human ovarian carcinoma. Mutations in this gene cause myopathy and distal arthrogryposis type 2B. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Mar 2009]

Synonyms:

AMCD2B; DA2B; FSSV; fsTnl

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TNNI2 ([RC205676], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TNNI2. Positive lysates [LY418789] (100ug) and [LC418789] (20ug) can be purchased separately from OriGene.