

Product datasheet for CF808066

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TNFRSF14 Mouse Monoclonal Antibody [Clone ID: OTI7D7]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7D7

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human Host: Mouse

Isotype: lgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 39-283 of human

TNFRSF14(NP_003811) produced in E.coli.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 26.1 kDa

Gene Name: TNF receptor superfamily member 14

Database Link: NP 003811

Entrez Gene 8764 Human

Q92956



TNFRSF14 Mouse Monoclonal Antibody [Clone ID: OTI7D7] - CF808066

Background: This gene encodes a member of the TNF (tumor necrosis factor) receptor superfamily. The

encoded protein functions in signal transduction pathways that activate inflammatory and inhibitory T-cell immune response. It binds herpes simplex virus (HSV) viral envelope glycoprotein D (gD), mediating its entry into cells. Alternative splicing results in multiple


transcript variants. [provided by RefSeq, Jul 2014]

Synonyms: ATAR; CD270; HVEA; HVEM; LIGHTR; TR2

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Cytokine-cytokine receptor interaction

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TNFRSF14 ([RC201167], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TNFRSF14 (1:2000). Positive lysates [LY401254] (100ug) and [LC401254] (20ug) can be purchased separately from OriGene.