

## **Product datasheet for CF808308**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# NOTCH1 Mouse Monoclonal Antibody [Clone ID: OTI2E7]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2E7
Applications: IHC

Recommended Dilution: IHC 1:150

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 2280-2556 of human

NOTCH1(NP\_060087) produced in E.coli.

**Formulation:** Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 270.6 kDa

**Gene Name:** notch receptor 1

Database Link: NP 060087

Entrez Gene 18128 MouseEntrez Gene 25496 RatEntrez Gene 4851 Human

P46531





Background:

This gene encodes a member of the Notch family. Members of this Type 1 transmembrane protein family share structural characteristics including an extracellular domain consisting of multiple epidermal growth factor-like (EGF) repeats, and an intracellular domain consisting of multiple, different domain types. Notch family members play a role in a variety of developmental processes by controlling cell fate decisions. The Notch signaling network is an evolutionarily conserved intercellular signaling pathway which regulates interactions between physically adjacent cells. In Drosophilia, notch interaction with its cell-bound ligands (delta, serrate) establishes an intercellular signaling pathway that plays a key role in development. Homologues of the notch-ligands have also been identified in human, but precise interactions between these ligands and the human notch homologues remain to be determined. This protein is cleaved in the trans-Golgi network, and presented on the cell surface as a heterodimer. This protein functions as a receptor for membrane bound ligands, and may play multiple roles during development. [provided by RefSeq, Jul 2008]


**Synonyms:** AOS5; AOVD1; hN1; TAN1

Protein Families: Adult stem cells, Cancer stem cells, Druggable Genome, ES Cell Differentiation/IPS, Stem cell

relevant signaling - DSL/Notch pathway

**Protein Pathways:** Dorso-ventral axis formation, Notch signaling pathway, Prion diseases

# **Product images:**



Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-NOTCH1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.