

# **Product datasheet for CF808996**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PD L2 (PDCD1LG2) Mouse Monoclonal Antibody [Clone ID: OTI6H10]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI6H10
Applications: FC, WB
Recommended Dilution: WB 1:500
Reactivity: Human
Host: Mouse

**Isotype:** IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PDCD1LG2 (NP\_079515) produced in

HEK293T cell.

**Formulation:** Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 30.8 kDa

**Gene Name:** programmed cell death 1 ligand 2

Database Link: NP 079515

Entrez Gene 80380 Human

Q9BQ51

**Background:** Involved in the costimulatory signal, essential for T-cell proliferation and IFNG production in a

PDCD1-independent manner. Interaction with PDCD1 inhibits T-cell proliferation by blocking

cell cycle progression and cytokine production. [UniProtKB/Swiss-Prot Function]






Synonyms: B7DC; bA574F11.2; Btdc; CD273; PD-L2; PDCD1L2; PDL2


**Protein Families:** Transmembrane

**Protein Pathways:** Cell adhesion molecules (CAMs)

### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PDCD1LG2 ([RC224141], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDCD1LG2 (1:500). Positive lysates [LY410802] (100ug) and [LC410802] (20ug) can be purchased separately from OriGene.



Flow cytometric analysis of living 293T cells transfected with PDL2 overexpression plasmid ([RC224141], Red)/empty vector ([PS100001], Blue) using anti-PDL2 antibody ([TA808996]). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).