

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for CF810618

BAG5 Mouse Monoclonal Antibody [Clone ID: OTI3C10]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI3C10
Applications:	WB
Recommended Dilution:	WB 1:500~2000
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human BAG5 (NP_004864) produced in 293T.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	BAG cochaperone 5
Database Link:	<u>NP_004864</u> <u>Entrez Gene 70369 MouseEntrez Gene 366734 RatEntrez Gene 9529 Human</u> <u>Q9UL15</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


SorigeneBAG5 Mouse Monoclonal Antibody [Clone ID: OTI3C10] - CF810618Background:The protein encoded by this gene is a member of the BAG1-related protein family. BAG1 is an
anti-apoptotic protein that functions through interactions with a variety of cell apoptosis and
growth related proteins including BCL-2, Raf-protein kinase, steroid hormone receptors,
growth factor receptors and members of the heat shock protein 70 kDa family. This protein
contains a BAG domain near the C-terminus, which could bind and inhibit the chaperone
activity of Hsc70/Hsp70. Three transcript variants encoding two different isoforms have been
found for this gene. [provided by RefSeq, Jul 2008]

Synonyms:	BAG-5
Protein Families:	Druggable Genome

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BAG5 ([RC208518], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BAG5 (1:2000). Positive lysates [LY417691] (100ug) and [LC417691] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 7 different cell lines by using anti-BAG5 monoclonal antibody (1:500).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US