

Product datasheet for CF811180

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

AIF (AIFM1) Mouse Monoclonal Antibody [Clone ID: OTI8B8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI8B8

Applications: WB

Recommended Dilution: WB 1:500~2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 134-483 of human

AIFM1 (NP 001124318) produced in E.coli.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: apoptosis inducing factor mitochondria associated 1

Database Link: NP 001124318

Entrez Gene 26926 MouseEntrez Gene 83533 RatEntrez Gene 9131 Human

<u>095831</u>

Synonyms: AIF; CMT2D; CMTX4; COWCK; COXPD6; NADMR; NAMSD; PDCD8


Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Apoptosis






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY AIFM1 (Cat# [RC224064], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-AIFM1 (Cat# [TA811180])(1:2000).



Western blot analysis of extracts (35ug) from 6 different cell lines and mouse liver tissue lysate by using anti-AIFM1 monoclonal antibody (1:500).





Equivalent amounts of cell lysates (10 ug per lane) of wild-type 293T cells (WT, Cat# LC810293T) and AIFM1-Knockout 293T cells (KO, Cat# [LC811853]) were separated by SDS-PAGE and immunoblotted with anti-AIFM1 monoclonal antibody [TA811180], (1:500). Then the blotted membrane was stripped and reprobed with antib-actin antibody ([TA811000]) as a loading control.