

Product datasheet for CF812184

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

DNAH6 Mouse Monoclonal Antibody [Clone ID: OTI5C2]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5C2
Applications: IHC, WB

Recommended Dilution: WB 1:500, IHC 1:500

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human DNHL1 (NP_775916) produced in E.coli.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 46.6 kDa

Gene Name: dynein axonemal heavy chain 6

Database Link: NP 775916

Entrez Gene 284944 Entrez Gene 1768 Human


Q9C0G6

Synonyms: FLJ37357






Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY DNHL1 (Cat# [RC211192], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DNHL1(Cat# [TA812184]). Positive lysates [LY406530] (100ug) and [LC406530] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human placenta tissue within the normal limits using anti-DNHL1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA812184]) (1:500)