

Product datasheet for CF812307

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TTC7A Mouse Monoclonal Antibody [Clone ID: OTI4A2]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4A2

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-450 of human TTC7A

(NP_065191) produced in E.coli.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 96 kDa

Gene Name: tetratricopeptide repeat domain 7A

Database Link: NP 065191

Entrez Gene 362696 RatEntrez Gene 57217 Human

Q9ULT0





TTC7A Mouse Monoclonal Antibody [Clone ID: OTI4A2] - CF812307


Background:

This gene encodes a protein containing tetratricopeptide repeats. Mutations in this gene disrupt intestinal development and can cause early onset inflammatory bowel disease and intestinal atresia. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2014]

Synonyms:

GIDID; MINAT; TTC7

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TTC7A ([RC217177], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TTC7A. Positive lysates [LY412463] (100ug) and [LC412463] (20ug) can be purchased separately from OriGene.