

# **Product datasheet for CF812548**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## beta Arrestin 1 (ARRB1) Mouse Monoclonal Antibody [Clone ID: OTI3G9]

#### **Product data:**

**Applications:** 

**Product Type:** Primary Antibodies

Clone Name: OTI3G9

Recommended Dilution: WB 1:500

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ARRB1 (NP\_004032) produced in E.coli.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if

necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 47.1 kDa

**Gene Name:** arrestin beta 1

Database Link: NP 004032

Entrez Gene 25387 RatEntrez Gene 109689 MouseEntrez Gene 408 Human

P49407





Background:


Members of arrestin/beta-arrestin protein family are thought to participate in agonist-mediated desensitization of G-protein-coupled receptors and cause specific dampening of cellular responses to stimuli such as hormones, neurotransmitters, or sensory signals. Arrestin beta 1 is a cytosolic protein and acts as a cofactor in the beta-adrenergic receptor kinase (BARK) mediated desensitization of beta-adrenergic receptors. Besides the central nervous system, it is expressed at high levels in peripheral blood leukocytes, and thus the BARK/beta-arrestin system is believed to play a major role in regulating receptor-mediated immune functions. Alternatively spliced transcripts encoding different isoforms of arrestin beta 1 have been described. [provided by RefSeq, Jan 2011].

Synonyms: ARB1; ARR1

**Protein Families:** Druggable Genome

**Protein Pathways:** Chemokine signaling pathway, Endocytosis, MAPK signaling pathway

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY ARRB1 (Cat# [RC201279], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ARRB1 antibody (Cat# [TA812548])(1:500)