

Product datasheet for CF812580

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD272 (BTLA) Mouse Monoclonal Antibody [Clone ID: OTI7G4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7G4

Applications: FC

Recommended Dilution: FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 31-157+Mouse Fc of

human BTLA (NP_861445) produced in HEK293T cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 32.6 kDa

Gene Name: B and T lymphocyte associated

Database Link: NP 861445

Entrez Gene 151888 Human

Q7Z6A9






Background: This gene encodes a member of the immunoglobulin superfamily. The encoded protein

contains a single immunoglobulin (Ig) domain and is a receptor that relays inhibitory signals to suppress the immune response. Alternative splicing results in multiple transcript variants. Polymorphisms in this gene have been associated with an increased risk of rheumatoid

arthritis. [provided by RefSeq, Aug 2011]

Synonyms: BTLA1; CD272
Protein Families: Transmembrane

Product images:



Flow cytometric analysis of living 293T cells transfected with BTLA overexpression plasmid ([RC219458]), Red)/empty vector ([PS100001], Blue) using anti-BTLA antibody ([TA812580]). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).