

Product datasheet for CF813033

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

TIGIT Mouse Monoclonal Antibody [Clone ID: OTI1E6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1E6
Applications: FC, WB

Recommended Dilution: WB 1:1000, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TIGIT (NP_776160) produced in HEK293T

cell

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 26.1 kDa

Gene Name: T cell immunoreceptor with Ig and ITIM domains

Database Link: NP 776160

Entrez Gene 201633 Human

Q495A1





Background: This gene encodes a member of the PVR (poliovirus receptor) family of immunoglobin


proteins. The product of this gene is expressed on several classes of T cells including follicular B helper T cells (TFH). The protein has been shown to bind PVR with high affinity; this binding is thought to assist interactions between TFH and dendritic cells to regulate T cell dependent

B cell responses. [provided by RefSeq, Sep 2009]


Synonyms: VSIG9; VSTM3; WUCAM

Protein Families: Transmembrane

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY TIGIT (Cat# [RC221447], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TIGIT (Cat# [TA813033])(1:1000).



Flow cytometric analysis of living 293T cells transfected with TIGIT overexpression plasmid ([RC221447]), Red)/empty vector ([PS100001], Blue) using anti-TIGIT antibody ([TA813033]). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).