

Product datasheet for CF813093

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD137 (TNFRSF9) Mouse Monoclonal Antibody [Clone ID: OTI7C3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7C3

Applications: FC

Recommended Dilution: FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TNFRSF9 (NP_001552) produced in

HEK293T cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: TNF receptor superfamily member 9

Database Link: NP 001552

Entrez Gene 3604 Human

Q07011

Background: Receptor for TNFSF9/4-1BBL. Possibly active during T cell activation. [UniProtKB/Swiss-Prot

Function]


Synonyms: 4-1BB; CD137; CDw137; ILA







Protein Families: Druggable Genome, Transmembrane
Protein Pathways: Cytokine-cytokine receptor interaction

Product images:





Flow cytometric analysis of living 293T cells transfected with TNFRSF9 overexpression plasmid ([RC200664]), Red)/empty vector ([PS100001], Blue) using anti-TNFRSF9 antibody ([TA813093]). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).





Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h (Right)/untreated (Left) using anti-TNFRSF9 antibody ([TA813093]) (1:100).





Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h (Right) using anti-TNFRSF9 antibody ([TA813093]). Cells incubated with a non-specific antibody (Left) were used as isotype control (1:100).