

# **Product datasheet for CF813575**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CD133 (PROM1) Mouse Monoclonal Antibody [Clone ID: OTI5B12]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI5B12
Applications: FC, WB

Recommended Dilution: WB 1:1000, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PROM1 (NP\_006008) produced in HEK293T

cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if

necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 97 kDa

Gene Name: prominin 1

Database Link: NP 006008

Entrez Gene 8842 Human

043490






#### Background:

This gene encodes a pentaspan transmembrane glycoprotein. The protein localizes to membrane protrusions and is often expressed on adult stem cells, where it is thought to function in maintaining stem cell properties by suppressing differentiation. Mutations in this gene have been shown to result in retinitis pigmentosa and Stargardt disease. Expression of this gene is also associated with several types of cancer. This gene is expressed from at least five alternative promoters that are expressed in a tissue-dependent manner. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2009].


Synonyms: AC133; CD133; CORD12; MCDR2; MSTP061; PROML1; RP41; STGD4

**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Transmembrane

# **Product images:**





HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PROM1 ([RC221611], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PROM1.(1:1000)




Western blot analysis of extracts (35ug) from 4 cell lines lysates by using anti-PROM1 monoclonal antibody. (1:1000)











Flow cytometric analysis of living HT29 cells, using anti-PROM1 antibody([TA813575], Red), compared to an isotype control (green), and a PBS control (blue).(1:100)



Flow cytometric analysis of living HCT116 cells, using anti-PROM1 antibody([TA813575], Red), compared to an isotype control (green), and a PBS control (blue).(1:100)





Flow cytometric analysis of living NCCIT cells, using anti-PROM1 antibody([TA813575], Red), compared to an isotype control (green), and a PBS control (blue).(1:100)